



Area Size

152 890 km²

Qualifying Species and Criteria

Amazon river dolphin – *Inia geoffrensis*

Criterion A; C (2)

West Indian manatee – *Trichechus manatus*

Criterion A; C (1, 2)

Amazonian manatee – *Trichechus inunguis*

Criterion A; C (1); D1

Guiana dolphin – *Sotalia guianensis*

Criterion B (1); C (1)

Marine Mammal Diversity

Criterion D (2)

Inia geoffrensis, *Trichechus manatus*, *Trichechus inunguis*, *Sotalia guianensis*, *Tursiops truncatus*, *Megaptera novaeangliae*, *Steno bredanensis*, *Globicephala macrorhynchus*, *Pseudorca crassidens*, *Lagenodelphis hosei*, *Delphinus delphis*, *Grampus griseus*, *Stenella longirostris*, *Pteronura brasiliensis*, *Lontra longicaudis*

Summary

The Guianas to Amazon Outflow IMMA extends from central Guianas to Maranhão, Brazil overlapping with the Amazonian-Orinoco Influence Zone EBSA. A predominant feature of the IMMA is the outflow, the world's largest river discharge, a low salinity (<35 ppt) plume with high

Guianas to Amazon Outflow IMMA

Summary, continued.

concentrations of suspended material and nutrients spreading over thousands of kilometers along the North Brazilian Continental Shelf and the Equatorial Atlantic Ocean. This freshwater plume and its associated nutrients lead to increased productivity, creating an important feeding area for the survival of a rich local community of aquatic mammals. The composition of the IMMA's cetacean community is more similar to those in the Caribbean than to cetaceans of the north-eastern Brazilian coast. The IMMA also encompasses the Great Amazon Reef System, which extends from French Guiana to Pará in Brazil. The IMMA provides habitat for several estuarine and coastal species that are threatened due to their restricted habitat requirements that overlap with human activity. These include the Guiana dolphin (*Sotalia guianensis*), the Amazon River dolphin (*Inia geoffrensis*). These dolphins rely on mangrove ecosystems and estuaries, forming discrete populations with small home ranges. Additionally, the IMMA provides habitat for both the Amazonian manatee (*Trichechus inunguis*) and the West Indian manatee (*Trichechus manatus*), representing unique contact zone where hybrids between both species occur.

Description:

This IMMA extends along the coasts of Suriname (up to the Suriname River), French Guiana and northern Brazil (up to São Luis, Maranhão state), and is part of the Amazonian-Orinoco Influence Zone EBSA. Within this IMMA are eight Ramsar sites, including the Amazon estuary and its mangroves.



Figure 1: Guiana dolphins (*Sotalia guianensis*). Photo credit: Collection of Green Heritage Fund Suriname

The estuary of the Amazon and its mangroves cover 3.8 million hectares in Amapá, Pará and Maranhão states, including 23 conservation units primarily designated as sustainable use protected areas. In the east of the IMMA, the Marajoara and Maranhense gulfs in Maranhão state represent dynamic estuarine complexes facilitating substantial solid discharges. Estuaries, coastal lagoons and mangroves are present along the entire northern coast (Abreu, 2015).

The coastal and marine habitats feature shallow depths over extensive mudflats along mostly intact coastal mangrove forests of *Avicennia germinans*, which in French Guiana are interspersed with rocky outcrops. The gently sloping continental shelf extends up to 150 km offshore in French Guiana and includes the Great Amazon Reef System (GARS).

The IMMA is an important environment for two manatee species, the Amazonian manatee (*Trichechus inunguis*) and the Antillean subspecies of the West Indian manatee (*Trichechus manatus manatus*), and two dolphin species, the Guiana dolphin (*Sotalia guianensis*) and the Amazon River dolphin (*Inia geoffrensis*) and its Araguaian population (Siciliano et al., 2016; Costa et al., 2017).

Criterion A: Species or Population Vulnerability

The Guiana dolphin (*Sotalia guianensis*) has a discontinuous distribution along most of its range along the Atlantic coast of northern and eastern South America, and small resident populations can be found in distinct pockets of habitat in mangrove



Figure 2: *Inia araguaiaensis* in Mocajuba. Photo credit: Paulo H. Ott

ecosystems, estuaries and bays (Bordin et al., 2022; Domit et al., 2021; Marmontel et al., 2021; Pool, 2019; Domit et al., 2021). Although the species is thought to occur along this entire range, Guiana dolphins usually form discrete populations, in which individuals typically have relatively small home ranges (Domit et al., 2021 and references therein). In 2018, the global IUCN Red list status of the Guiana dolphin was changed from Data Deficient to Near Threatened (Secchi et al., 2018). However, in French Guiana the Guiana dolphin is listed as *Endangered* (Moncorps et al., 2017) and in Brazil as *Vulnerable* (Secchi et al., 2018).

The Amazon Outflow IMMA is important for the survival and recovery of the Amazon River dolphin (*Inia geoffrensis*) which has been assessed as *Endangered* on the IUCN Red List of Threatened

Species (Da Silva et al., 2018), and two species of manatees: the West Indian (*Trichechus manatus*) and the Amazonian manatee (*T. inunguis*) both globally classified as *Vulnerable* on the IUCN Red List (Deutsch et al., 2008; Marmontel et al., 2016) and *Endangered* in French Guiana (Moncorps et al., 2017). Moreover, the Amazon River plume has led to the expansion of the geographic distribution of the Araguaian form of *Inia geoffrensis*, the only endemic cetacean of Brazil, into the shoreline of Marajó Bay and to the east, into the Curuçá Estuary, the eastern part of the Amazon Delta (Pará River) and approximately 500 km downstream of the Tucurui Dam (Siciliano et al., 2016).

Some studies indicate that the West Indian manatee population in South America has declined over the years due to various anthropogenic factors (Lima et



Figure 3: Dolphin perfume charm" from Tijuca, Rio de Janeiro, Rio de Janeiro state. Usage of cetacean product as 'love charm' in Brazil had threatened the Amazon river dolphin population. Photo credit: Siciliano et al. (2018)

al., 2011; Alves et al., 2013; Alves et al., 2015). Meirelles et al. (2022) evaluated the extinction risk of the West Indian manatee in Brazil. Although available data on the species' abundance and mortality are not sufficient to apply the IUCN criteria, the authors worked with limited information following the guidelines, inferring, and projecting based on evidence to produce information and properly evaluate the West Indian manatee risk of extinction. The results indicated that *T. manatus* is Critically Endangered in Brazil due to the extremely high suspected and projected population reduction ($\geq 80\%$) in the past and the future. Furthermore, although the causes of significant mortality were reduced in some parts of the species' range, such as from hunting, other threats are increasing and are not expected to decrease in the coming decades.

Bonvicino (2020) established the presence of the Amazon manatee throughout the lower Amazon and

its tributaries, including Marajó Island and its Atlantic coast and coastal islands, and from the northern (Amapá state, Brazil) to the southern shore of the Amazon estuaries (Pará state, Brazil). The current IMMA also hosts a hybridization zone of West Indian and Amazonian manatees. There are records of interspecific hybrids of *T. inunguis* and *T. manatus* close to the Amazon River mouth and along the Guiana Shield coastline (Noronha et al., 2022).

Criterion B: Distribution and Abundance

Sub-Criterion B1: Small and Resident Populations

The Guiana dolphin has a discontinuous distribution along most of its range and small resident populations are present year-round in small pockets in estuaries and bays (Bordin et al., 2022; Marmontel et al., 2021; Pool, 2019). These areas are used by these resident populations for feeding, mating, resting, and



Figure 4: Guiana dolphins (*Sotalia guianensis*) spyhopping. Photo credit: Marijke N. de Boer

all biological functions related to reproduction, such as giving birth, nursing and caring for calves (Pool, 2019).

There are few abundance/density studies of Guiana dolphin distribution. Large scale aerial surveys conducted in 2008 and 2017 estimated Guiana dolphin abundance/density in French Guiana to be roughly 1,700 individuals (95% CI: 700-4 000; Laran et al., 2019; Mannocci et al., 2013). However, with coastal aerial surveys conducted in 2013, 2014 and 2019 concluded lower estimates (Bordin et al., 2022).

Currently, the size of the population of Guiana dolphins in the Suriname River estuary is estimated to be between 80-100 based on photo identification. Of these, 67 are identified using dorsal fin identification and mark and recapture methodology, and some individuals have been resighted on multiple occasions since 2007 (Pool, 2019).

Criterion C: Key Life Cycle Activities

Sub-Criterion C1: Reproductive Areas

The West Indian manatee and the Amazonian manatee occur in the Marajó Bay area and in the inlets and channels near Belém (under the influence of the Amazon River Plume). The Amazon Aquatic Mammal Study Group (GEMAM) coordinates a collaborative network to locate manatees trapped in fishing nets and conduct rescues along the coast of the state of Pará and in the interior of Belém. The presence of the Amazonian manatee is confirmed with the rescue and sightings of calves and adults in the great Belém area (Emin-Lima et al., 2021). On October 31st 2021, Amazonian manatees were sighted swimming near a beach in the Mosqueiro district, in greater Belém where mating behaviour was filmed by residents and tourists.

Both species of manatees are threatened by anthropogenic factors, such as habitat loss due to coastal mangrove destruction, and bycatch in fishing gear (Parente et al., 2004; Borges et al., 2007; Meirelles, 2008; Silva et al., 2016).

Strandings of neonates and calves of both species have been reported in the Amazon estuary, Tocantins Bay and Oiapoque estuary, all important breeding sites for both, West Indian and Amazonian manatees (Parente et al., 2004; Borges et al., 2007; Meirelles, 2008; Silva 2015).

Sub-Criterion C2: Feeding Areas

The discharge of the Amazon River plume promotes a change in the freshwater layer and consequently in local productivity (Tosseto et al., 2022), creating an important feeding area for all coastal species, including Guiana dolphins and Amazonian manatees, as well as other aquatic vertebrates, such as fishes and sea birds (Siciliano et al., 2008; Costa et al., 2017; Costa et al., 2020; Tosseto et al., 2022).

The coastal waters of the IMMA are largely influenced by the massive suspended-sediment discharge, causing mega-turbidity on the shelf, a growth of a subaqueous delta and an intermediary salinity (Tosseto et al., 2022). All these factors have a direct influence on oceanic biodiversity, including marine species, coral reef endemics and threatened species completely dependent on ecosystems such as estuaries, mudflats, sandy beaches and, particularly, littoral mangroves (Artigas et al., 2003).

A study of bone and teeth samples collected from museum samples and stranded or floating aquatic mammal carcasses in the Amazon estuarine complex and adjacent coastal zone and along Northeastern coast of Brazil included 220 samples of Guiana dolphins, and 11 from Amazonian manatees. Stable

isotope analysis to analyse trophic relationships revealed that both species were feeding in the area, and that there was clear resource partitioning between Guiana dolphins and Amazonian manatees. Isotopic niches were highly distinct, with no overlap among these two species inhabiting the Amazon plume. The study also indicated that both species had a fairly wide variety in their prey base and their 'isotopic niche', suggesting high habitat plasticity (Costa et al., 2020).

Criterion D: Special Attributes

Sub-Criterion D1: Distinctiveness

The West Indian (*Trichechus manatus*) and Amazonian (*T. inunguis*) manatee have an overlapping distribution at the mouth of the Amazon River. A result of this sympatry is the occurrence of hybrids, which are frequently found along the coasts of Amapá state in Brazil, French Guiana and Guyana. New genetic evidence (Lima et al., 2019) indicates the occurrence of a concentration of hybrids along the Guianas Shield coastline, which is an interspecific hybrid zone that also separates *T. manatus* populations located east (Brazil) and west (Caribbean, Gulf of Mexico, Florida and Antilles). In addition, Lima et al. (2019) suggest that this hybrid population, which occupies a unique mangrove-rich environment under strong influence of the Amazon River plume, requires independent conservation management measures.

The presence of hybrid manatees in this area under influence of the Amazon plume seems to be correlated both to the isolation of the western and eastern *Trichechus m. manatus* and the sympatric area between *T. inunguis* and *T. m. manatus* specimens (Vianna et al., 2006; Santos et al., 2016; de Oliveira et al., 2022).



Figure 5: *Omar* the hybrid manatee from Marajó Island, Pará. Photo credit: R. Baleia

Sub-Criterion D2: Diversity

The Amazon Outflow encompasses the Amazon River plume, including estuarine, coastal and oceanic zones. According to Siciliano et al. (2008) and Costa et al. (2017) local aquatic mammal richness ranged from 15 to 22 species, and the most common species are Guiana dolphins, Amazon river dolphins, and West Indian manatees (Siciliano et al., 2008; Santos et al., 2016; Costa et al., 2017). Common bottlenose dolphins (*Tursiops truncatus*) are also common throughout the IMMA (Mannocci, 2013; Pelagis, 2021). Additional species encountered during vessel-based and aerial surveys of the coastal areas of the IMMA include humpback whales (*Megaptera novaeangliae*), short-finned pilot whales (*Globicephala macrorhynchus*), false killer whales (*Pseudorca*

crassidens), Risso's dolphins (*Grampus griseus*), rough-toothed dolphins (*Steno bredanensis*), Fraser's dolphins (*Lagenodelphis hosei*), common dolphins (*Delphinus delphis*), and spinner dolphins (*Stenella longirostris*) (Ramos et al., 2001; Costa et al., 2017; Pelagis, 2021; <https://www.obsenmer.org/maps>; Pusineri et al., 2021; De Boer, 2022). In addition, two species of otters are encountered in the area – giant otters (*Pteronura brasiliensis*) (e.g. Carter & Rosas, 2008; Duplaix et al., 2018) and neotropical otters (*Lontra longicaudis*) (e.g. Lopes Rheingantz, 2014).

Supporting Information

Abreu, C.T. 2015. Brazilian Coastal and Marine Protected Areas Importance, Current Status and Recommendations. University of Rhode Island.

Agência Nacional de Águas (Brasil) (ANA). 2015. HidroWeb: Sistemas de informações hidrológicas. Disponível em: <http://hidroweb.ana.gov.br/HidroWeb>. Accessed 22 January 2015.

Alves, M.D., Kinás, P.G., Marmontel, M., Borges, J.C.G., Costa, A.F., Schiel, N., and Arujo, M.E. 2015. First abundance estimate of the Antillean manatee (*Trichechus manatus manatus*) in Brazil by aerial survey. Journal of the Marine Biological Association of the United Kingdom, 1, 1–12.

Alves, M.D.O., Schwamborn, R., Borges, J.C.G., Marmontel, M., Costa, A.F., Schettini, C.A.F., Araújo, M.E. 2013. Aerial survey of manatees, dolphins and sea turtles off northeastern Brazil: Correlations with coastal features and human activities. Biol. Conserv. 161, 91–100. doi:10.1016/j.biocon.2013.02.015

Artigas, L.F., Vendeville, P., Leopold, M., Guiral, D., and Ternon, J.F. 2003. Marine biodiversity in French Guiana: estuarine, coastal, and shelf ecosystems under the influence of Amazonian waters. Gayana 67: 302–326.

Balensiefer, D.C., Attademo, F.L.N., Sousa, G.P., Freire, A.C. da B., da Cunha, F.A.G.C., Alencar, A.E.B., Silva, F.J. de L., Luna, F., and de O. 2017. Three Decades of Antillean Manatee (*Trichechus manatus manatus*) Stranding Along the Brazilian Coast. Tropical Conservation Science 10, 194008291772837. <https://doi.org/10.1177/1940082917728375>.

Barroso, C.X., Lotufo, T.M.C. and Matthews-Cascon, H. 2016. Biogeography of Brazilian prosobranch gastropods and their Atlantic relationships. J. Biogeogr. 43, 2477–2488.

Bezerra, M.O., Medeiros, C., Krelling, A.P.M., Rosário, R.P., and Rollnic, M. 2011. Physical oceanographic

behavior at the Guamá/Acará-Mojú and the Paracauari river mouths, Amazon coast (Brazil). In: Proceedings of the 11th international coastal symposium, Szczecin. Journal of Coastal Research 2011: 1448–1452.

Bonvicino, C.R., Viana, M.C., Oliveira, E.H.C. de, Emin, R., Silva Junior, J.D.S. e, Sousa, M.E.M. de, Siciliano, S., 2020. Distribution of South American manatees, *Trichechus manatus* Linnaeus, 1758 and *T. inunguis* (Natterer, 1883) (Sirenia: Trichechidae). bcnaturais 15, 573–599.

<https://doi.org/10.46357/bcnaturais.v15i3.246>.

Bordin, A., Vanhoucke, M., Pineau, K., Kelle, L., Cozannet, N., Pool, M., Bolaños-Jiménez, J., and De Thoisy, B. 2022. Study and conservation of the Guiana Dolphin (*Sotalia guianensis*) (Van Bénéden, 1864) in French Guiana. Lat. Am. J. Aquat. Mamm. <https://doi.org/10.5597/lajamoo276>.

Borges, J.C.G., Vergara-Parente, J.E., Alvite, C.M. de C., Marcondes, M.C.C., Lima, R.P. de, 2007. Embarcações motorizadas: uma ameaça aos peixes-boi marinhos (*Trichechus manatus*) no Brasil. Biota Neotrop. 7, 199–204. <https://doi.org/10.1590/S1676-06032007000300021>.

Borges, J.C.G., Araújo, P.G., Anzolin, D.G., and Miranda, G.E.C. 2008. Identificação de itens alimentares constituintes da dieta dos peixes-boi marinhos (*Trichechus manatus*) na região nordeste do Brasil. Biotemas. 21 (2), 77–81.

Brandini, F.P., Lopes, R.M., Gutseit, K.S., and Sassi, R. 2007. Planctonologia na Plataforma Continental do Brasil: Diagnose e Revisão Bibliográfica. (CEMAR/MMA/CIRM/FEMAR, 1997).

Carter, S.K. and Rosas, F.C.W. 1997. Biology and conservation of the Giant Otter *Pteronura brasiliensis*.

- Mammal Review, 27: 1-26.
<https://doi.org/10.1111/j.1365-2907.1997.tb00370.x>.
- Chollett, I., Mumby, P.J., Müller-Karger, F.E., and Hu, C. 2012. Physical environments of the Caribbean Sea. Limnol. Oceanogr. 57: 1233-1244.
- Costa, M.S. 2013. Aporte hídrico e do material particulado em suspensão para a baía do Marajó: influência dos rios Amazonas e Tocantins. Master Thesis, Universidade Federal do Pará.
- Costa, A.F., Siciliano, S., Emin-Lima, R., Martins, B.M.L., Sousa, M.E.M., Giarrizzo, T., and Silva Júnior, J.S. 2017. Stranding survey as a framework to investigate rare cetacean records of the north and north-eastern Brazilian coasts. ZooKeys 688: 111-134.
<https://doi.org/10.3897/zookeys.688.12636>.
- Costa, A.F., Botta, S., Siciliano, S. et al. 2020. Resource partitioning among stranded aquatic mammals from Amazon and Northeastern coast of Brazil revealed through carbon and nitrogen stable isotopes. Sci Rep 10, 12897. <https://doi.org/10.1038/s41598-020-69516-8>.
- da Silva, V., Trujillo, F., Martin, A., Zerbini, A.N., Crespo, E., Aliaga-Rossel, E., and Reeves, R. 2018. *Inia geoffrensis*. The IUCN Red List of Threatened Species 2018: e.T10831A50358152.
<https://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T10831A50358152.en>. Accessed on 07 June 2023.
- Duplaix, N., Evangelista, E. and Rosas, F.C.W. 2015. Advances in the study of giant otter (*Pteronura brasiliensis*) ecology, behavior, and conservation: a review. Latin American Journal of Aquatic Mammals 10(2): 75-98.
<http://dx.doi.org/10.5597/lajam00200>.
- de Mahiques, M.M., Siegle, E., Francini-Filho, R.B., Thompson, F.L., de Rezende, C.E., Gomes, J.D., and Asp, N.E. 2019. Insights on the evolution of the living Great Amazon Reef System, equatorial West Atlantic. Sci Rep 9, 13699. <https://doi.org/10.1038/s41598-019-50245-6>.
- de Oliveira EHC, Gomes AJB, Costa AF, Emin-Lima R, Bonvicino CR, Viana MC, Reis LMA, Vidal MD, Cavalcanti MVG, Attademo FLN, Luna FO, Siciliano S. 2022. Karyotypical confirmation of natural hybridization between two manatee species, *Trichechus manatus* and *Trichechus inunguis*. Life (Basel). 2022 Apr 20;12(5):616. doi: 10.3390/life12050616. PMID: 35629284; PMCID: PMC9145575.
- Deutsch, C.J., Self-Sullivan, C., and Mignucci-Giannoni, A. 2008. *Trichechus manatus*: The IUCN Red List of Threatened Species 2008: e.T22103A9356917 [Data set]. <https://doi.org/10.2305/IUCN.UK.2008.RLTS.T22103A9356917.en>. Accessed on 20 May 2023.
- Dos Santos, S.S., Medeiros, I.S., Rebelo, V.A., Carvalho, A.O.B., Dubut, J.P., Mantovani, J.E., Círiaco, R.D., Dos Santos, R.E.G., Marmontel, M., Normande, I.C., Veloso, T.M.G., and Borges, J.C.G. 2022. Home ranges of released West Indian manatees in Brazil. ORYX, 56 : 1-8.
- Domit, C., de Miranda, A.V., Azevedo, A., Costa, A.F., de Meirelles, C.O., Farro, A.P., Bracarense, A.P.L., Barreto, A.S., Zaccaron, S., Andriolo, A., Louzada, C.N., Bertozzi, C.P., Marcondes, D.S., Ruenes, G., Melo-Santos, G., Cunha, H., Maciel, I., Jr, J.L.B., Torres-Florez, J.P., Tullio, J.D., Groch, K.R., May-Collado, L., Vidal, L.G., Flach, L., Sánchez, L., Londoño, M.C.R., Cremer, M.J., Alonso, M.B., Laeta, M., Cantor, M., Marmontel, M., Espinoza-Rodríguez, N., Fruet, P., Hauser, R., Sousa-Lima, R., Paitach, R., Tardin, R.,

- Siciliano, S., Moura, S., Caballero, S., Bisi, T., Belleghem, T.V., Ridoux, V., Luz, V., Briceño, Y., Pendu, Y.L., 2021. Progress report of the *Sotalia guianensis* Intersessional Group: Status of the Current Knowledge and Action Plan 69.
- Emin-Lima, R., Costa, A.F., Attademo, F.L.N., Hauser-Davis, R.A., Luna, F., de O., and Siciliano, S. 2021. Amazonian Manatees (*Trichechus inunguis*) Inhabiting an Equatorial Metropolis: Historical Records and Mating Activity Near Belém, Northern Brazil - Sobre A Presença Do Peixe-Boi Amazônico (*Trichechus inunguis*) Próximo À Belém: Registros Históricos E Atividade Reprodutiva. Bol. Lab. Hidrobiol. 31. <https://doi.org/10.18764/1981-6421e2021.12>.
- Emin-Lima, N.E. et al. 2022. Morphological disparity in the skull of Amazon River dolphins of the genus *Inia* (Cetacea, Iniidae) is inconsistent with a single taxon. Journal of Mammalogy. gyaco39. <https://doi.org/10.1093/jmammal/gyaco39>.
- Figueiredo, A.G., Allison, M. and Nittrouer, C.A. 1991. Amazon discharge (Technical Report). Universidade Federal Fluminense, Niteroi.
- Francini-Filho, R.B. et al. 2018. Perspectives on the Great Amazon Reef: Extension, biodiversity, and threats. Front. Mar. Sci. 5, 142.
- Hrbek, T., da Silva, V.M.F., Dutra, N., Gravena, W., Martin, A.R., and Farias, I.P. 2014. A New Species of River Dolphin from Brazil or: How Little Do We Know Our Biodiversity. PLoS ONE 9, e83623. <https://doi.org/10.1371/journal.pone.0083623>.
- Jo, Y.H. 2005. A study of the freshwater discharge from the Amazon River into the tropical Atlantic using multi-sensor data. Geophys. Res. Lett. 32, L02605. <https://doi.org/10.1029/2004GL021840>.
- Laran, S., Bassols, N., Dorémus, G., Authier, M., Ridoux, V., and Van Canneyt, O. 2019. Distribution et abondance de la mégafaune marine aux Petites Antilles et en Guyane française. Campagne REMMOA - II. Rapport final pour l'Agence Française pour la Biodiversité. https://www.observatoire-pelagis.cnrs.fr/wp-content/uploads/2021/05/12-RAPPORT_REMMOA_ANTGUY_FINAL_2019.pdf
- Lentz, S.J. and Limeburner, R. 1995. The Amazon River Plume during AMASSEDS: Spatial characteristics and salinity variability. J. Geophys. Res. 100, 2355. <https://doi.org/10.1029/94JC01411>.
- Lima, C.S., Magalhães, R.F., Marmontel, M., Meirelles, A.C., Carvalho, V.L., Lavergne, A., Thoisy, B.D., and Santos, F.R. 2019. A hybrid swarm of manatees along the Guianas coastline, a peculiar environment under the influence of the Amazon River plume. An. Acad. Bras. Ciênc. 91, e20190325. <https://doi.org/10.1590/0001-3765201920190325>.
- Lima, R.P., Paludo, D., Soavinski, R.J., Silva, K.G., and Oliveira, E.M.A. 2011. Levantamento da distribuição, ocorrência e status de conservação do Peixe-Boi Marinho (*Trichechus manatus*, Linnaeus, 1758) no litoral nordeste do Brasil. Natural Resources, 1: 41-57.
- Loder, J.W., Boicourt, W.C. and Simpson, J.H. 1998. Western ocean boundary shelves coastal segment (W). Sea 11, 3–27.
- Lopes Rheingantz, M., Saraiva de Menezes, J.F. and de Thoisy, B. 2014. Defining Neotropical otter *Lontra longicaudis* distribution, conservation priorities, and ecological frontiers. Tropical Conservation Science Vol.7 (2): 214-229. Available online: www.tropicalconservationscience.org.
- Luna, F. de O., de Lima, R.P., and de Araújo, J.P. 2008. Status de conservação do peixe-boi marinho

(*Trichechus manatus manatus* Linnaeus, 1758) no Brasil 10.

Mannocci, L., Monestiez, P., Bolaños-Jiménez, J., Dorémus, G., Jeremie, S., Laran, S., Rinaldi, R., Van Canneyt, O., and Ridoux, V. 2013. Megavertebrate communities from two contrasting ecosystems in the western tropical Atlantic. *Journal of Marine Systems* 111–112, 208–222.

Marmontel, M., de Souza, D. and Kendall, S. 2016. *Trichechus inunguis*: The IUCN Red List of Threatened Species 2016: e.T22102A43793736 [Data set]. International Union for Conservation of Nature. <https://doi.org/10.2305/IUCN.UK.2016-2.RLTS.T22102A43793736.en>.

Marmontel, M., Lima, D. dos S., Funi, C., Santos, V.F. dos, and Oliveira-da-Costa, M., 2021. Unveiling the Conservation Status of *Inia* and *Sotalia* in the Brazilian Northeastern Amazon. *Aquat Mamm* 47, 376–393. <https://doi.org/10.1578/AM.47.4.2021.376>.

Martins, S.E.M. and Mendes, A.C. 2011. Caracterização de depósitos sedimentares recentes da porção superior da Baía de Marajó (margem leste do estuário do Rio Pará, Amazônia). *Pesquisas em Geociências* 38(2): 168–180.

Meirelles, A.C.O. 2008. Mortality of the Antillean manatee, *Trichechus manatus manatus*, in Ceará State, northeastern Brazil. *Journal of the Marine Biological Association of the United Kingdom*, v. 88, n.6, p. 1133–1137.

Meirelles, A.C.O., Carvalho, V.L. and Marmontel, M. 2018. West Indian Manatee *Trichechus manatus* in South America: Distribution, Ecology and Health Assessment, in: Rossi-Santos, M.R., Finkl, C.W. (Eds.), *Advances in Marine Vertebrate Research in Latin America*, Coastal Research Library. Springer

International Publishing, Cham, pp. 263–291. https://doi.org/10.1007/978-3-319-56985-7_11.

Meirelles, A.C.O., Lima, D. dos S., Alves, M.D. de O., Borges, J.C.G., Marmontel, M., Carvalho, V.L., and Santos, F.R. dos. 2022. Don't let me down: West Indian manatee, *Trichechus manatus*, is still critically endangered in Brazil. *Journal for Nature Conservation* 67, 126169. <https://doi.org/10.1016/j.jnc.2022.126169>.

Moncorps, S., Siblet, J.P., Colas, H., Gigot, G., Jeusset, A., Kirchner, F., Allard, L., Brosse, S., Covain, R., Gozlan, R., Le Bail, P.Y., Melki, F., Vigouroux, R., Anselin, A., Delvaux, H., Rey, S., Villettard, M., and Virevaise, M. 2017. La liste rouge des espèces menacées en France : faune vertébrée de Guyane (No. fdi:o10072525). UICN, Paris.

Moura, R.L. et al. 2016. An extensive reef system at the Amazon River mouth. *Sci. Adv.* 2, 1–11.

Muehe, D. and Garcez, D.S. 2005. A Plataforma Continental Brasileira e Sua Relação com a Zona Costeira E A Pesca 20.

Neumann-Leitão, S. et al. 2018. Zooplankton from a reef system under the influence of the Amazon River plume. *Front. Microbiol.* 9, 1–15.

Nittrouer, C.A. and DeMaster, D.J. 1996. The Amazon shelf setting: tropical, energetic, and influenced by a large river. *Continental Shelf Research* 16, 553–573. [https://doi.org/10.1016/0278-4343\(95\)00069-0](https://doi.org/10.1016/0278-4343(95)00069-0).

Noronha, R.C.R., Almeida, B.R.R., Chagas, M.C.S., Tavares, F.S., Cardoso, A.L., Bastos, C.E.M.C., Silva, N.K.N., Klautau, A.G.C.M., Luna, F.O., Attademo, F.L.N., Lima, D.S., Sabioni, L.A., Sampaio, M.I.C., Oliveira, J.M., Nascimento, L.A.S.D., Martins, C., Vicari, M.R., Nagamachi, C.Y., and Pieczarka, J.C. 2022. Karyotypes of manatees: New insights into hybrid formation

(*Trichechus inunguis* × *Trichechus m. manatus*) in the Amazon Estuary. *Genes (Basel)*. 13(7):1263. doi: 10.3390/genes13071263. PMID: 35886048; PMCID: PMC9323068.

Normande, I.C., Malhado, A.C.M., Reid, J., Viana, P.C., Savaget, P.V.S., Correia, R.A., Luna, F.O., and Ladle, R.J. 2016. Post-release monitoring of Antillean manatees: an assessment of the Brazilian rehabilitation and release programme. *Animal Conservation*, 19: 235–246.

Oliveira, E.H.C., Gomes, A.J.B., Costa, A.F., Emin-Lima, R., Bonvicino, C.R., Viana, M.C., Reis, L.M.A., Vidal, M.D., Cavalcanti, M.V.G., Attademo, F.L.N., Luna, F.O., and Siciliano, S. 2022. Karyotypical confirmation of natural hybridization between two manatee species, *Trichechus manatus* and *Trichechus inunguis*. *Life* 12, 5: 616. <https://doi.org/10.3390/life12050616>.

Pansard, K.C.A., Gurgel, H. de C.B., Andrade, L.C. de A., and Yamamoto, M.E. 2011. Feeding ecology of the estuarine dolphin (*Sotalia guianensis*) on the coast of Rio Grande do Norte, Brazil. *Marine Mammal Science* 27, 673–687. <https://doi.org/10.1111/j.1748-7692.2010.00436.x>.

Parente, C.L., Vergara-Parente, J.E. and Lima, R.P. 2004. Strandings of Antillean manatees, *Trichechus manatus manatus*, in northeastern Brazil. *Lat. Am. J. Aquat. Mamm.* 3, 69–75. <https://doi.org/10.5597/lajam00050>.

Pelagis. 2021. Visual sightings of scientific surveys. <https://pelabox.univlr.fr/pelagis/PelaObs/> Accessed May 2023.

Pool, M.S. 2019. Estuarine Mammal Baseline for the proposed Suriname River Dredging ESIA Update Project.

Prestes, Y.O. 2013. Transporte de volume na zona de maré do rio Pará, Brasil. Monograph (Undergraduate), Institute of Geosciences, Universidade Federal do Pará.

Pusineri, C., Martinez, L., Bordin, A., Montgolfier, B., Deffes, O., Rufray, V., Lenrume, P., Safi, M., Suardi, A., Wyss, V., Geraldès, D., Marzelière, M., Leclerc, B., Barnicoat, S., and Rinaldi, R. 2021. Cetacean community off French Guiana. *Latin American Journal of Aquatic Mammals* 16(1): 12–22. <https://doi.org/10.5597/lajam00266>.

Ramos et al. 2001. A note on strandings and age of sperm whales (*Physeter macrocephalus*) on the Brazilian coast. *Journal of Cetacean Research and Management* 3(3):321–327.

Rocha, L.A. 2003. Patterns of distribution and processes of speciation in Brazilian reef fishes. *J. Biogeogr.* 30, 1161–1171.

Schwassmann, H.O., Barthem, R.B. and Carvalho, M.L. 1989. A note on the seasonally shifting zone of high primary production in the Bay of Marajó, Pará, Brazil, 1983–1984. *Acta Botanica Brasilica* 2(1): 165–174.

Secchi, E., Santos, M.C. de O. and Reeves, R. 2018. *Sotalia guianensis* (errata version published in 2019). The IUCN Red List of Threatened Species 2018: e.T181359A144232542. <https://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T181359A144232542.en>. Accessed on 25 January 2023.

Siciliano, S., Emin-lima, N.R., Costa, A.F., Tosi, C.H., Garri, R.G., Regina, C., Silva, D.A., Sousa, J.D.E., and Júnior, E.S. 2008. Revisão do conhecimento sobre os mamíferos aquáticos da Costa Norte do Brasil. *Arquivos do Museu Nacional* 66: 381–401.

Siciliano, S., Valiati, V.H., Emin-Lima, R., Costa, A.F.,

Sartor, J., Dorneles, T., de Sousa e Silva Júnior, J., and de Oliveira, L.R. 2016. New genetic data extend the range of river dolphins *Inia* in the Amazon Delta. *Hydrobiologia* 777, 255–269.
<https://doi.org/10.1007/s10750-016-2794-7>.

Siciliano, S. et al. 2023. Giving names to the characters: Identifying, tracing and estimating the multiple use of aquatic wildlife in Brazil. In: Galetti Jr., P.M. (eds) *Conservation Genetics in the Neotropics*. Springer, Cham. https://doi.org/10.1007/978-3-031-34854-9_13.

Silva, S.M. 2015. Caracterização populacional e estimativa de parentesco entre peixes-bois-marinhos *Trichechus manatus* (Linnaeus, 1758) usando marcadores microsatélites. Dissertação (Mestrado em Biologia Ambiental), Universidade Federal do Pará.

Silva, C.P.N., Meirelles, A.C.O. and Umezaki, J. 2016. Ameaças. In: Meirelles ACO, Carvalho, V.L. (eds) Peixe-boi-marinho: Biologia e Conservação no Brasil. São Paulo, Bambu, pp 93–108.

Targino, A.K.G. and Gomes, P.B. 2020. Distribution of sea anemones in the Southwest Atlantic: Biogeographical patterns and environmental drivers. *Mar. Biodivers.* 50, 80.

Tosetto G.E., Bertrand, A., Neumann-Leitão, S. et al. 2022. The Amazon River plume, a barrier to animal dispersal in the Western Tropical Atlantic. *Sci Rep* 12, 537. <https://doi.org/10.1038/s41598-021-04165-z>.

Acknowledgements

We would like to thank the participants of the 2023 hybrid IMMA Regional Expert Workshop for the identification of IMMAs in the South West Atlantic Ocean. Funding for the identification of this IMMA was provided by the Global Ocean Biodiversity Initiative funded by the German government's International Climate Initiative (IKI). Additional funds were provided by OceanCare and the Animal Welfare Institute. Local support and hosting for the workshop was provided by the Instituto Baleia Jubarte. Support to the IMMA programme is provided by Whale and Dolphin Conservation, and the Tethys Research Institute.



Suggested Citation: IUCN-MMPATF (2023) Guianas to Amazon Outflow IMMA Factsheet. IUCN Joint SSC/WCPA Marine Mammal Protected Areas Task Force, 2023.

PDF made available for download at
<https://www.marinemammalhabitat.org/portfolio-item/guianas-to-amazon-outflow-imma/>