


Madagascar Central East Coast IMMA

Description

The east coast of Madagascar is characterized by a very narrow continental shelf with a steep slope down to abyssal plain waters. In the central eastern portion, the continental shelf extends out to its widest point at approximately 5–60 km. The Masoala Peninsula in the northern part of this range defines Antongil Bay (16°00'S, 49°55'E), the largest bay in Madagascar, at approximately 60 km long and 30 km wide covering 2800 km². It is an expansive relatively shallow habitat with half being less than 50 m deep, and with 270 km of coastline. Île Sainte-Marie (16°55'S, 49°54'E), also known as Nosy Boraha, is an island off the east coast of Madagascar within the Analanjirofo administrative region. The island is 60 km long and less than 10 km wide and covers an area of 222 km². It is surrounded by a barrier reef that encloses a narrow, shallow-water lagoon (≤10 m depth). The Sainte Marie Channel (49°50' E–50°10' E, 16°60' S–17°55' S) lies between Île Sainte-Marie and the mainland of eastern Madagascar and is approximately 7 km wide with relatively shallow water depths (up to ~60 m).

The boundaries of the IMMA start in the north at the Masoala Peninsula (~15°S) and extend south to approximately 19.6°S, encompassing Antongil Bay, the waters off Île Sainte-Marie, the Sainte Marie Channel, and the narrow continental shelf past the port city of Toamasina (also known as Tamatave) to the mouth of the Mongoro River. The mouth of the Mongoro River (approx. 18°58' S, 49°50' E) demarcates the southern boundary of the IMMA. This area represents a bathymetric transition from a relatively shallow portion of the shelf that extends north to the Masoala Peninsula. The IMMA extends seaward approximately 50 nm, past the continental shelf edge.

Humpback whales are known to aggregate in the coastal waters of central eastern Madagascar during the winter, including in Antongil Bay, Île Sainte-Marie, and the waters south to the Mongoro River mouth (Rosenbaum et al., 1997; Best et al., 1998; Ersts and



Area Size

55,524 km²

Qualifying Species and Criteria

Humpback whale – *Megaptera novaeangliae*

Criteria B2, C1, C3

Marine Mammal Diversity

Feresa attenuata, *Indopacetus pacificus*,
Megaptera novaeangliae, *Peponocephala electra*,
Physeter macrocephalus, *Pseudorca crassidens*,
Tursiops aduncus, *Tursiops truncatus*, *Ziphius cavirostris*

Summary

The Madagascar Central East Coast IMMA encompasses Antongil Bay, the waters off Île Sainte-Marie, the narrow continental shelf past the port city of Toamasina (Tamatave) and waters beyond the continental shelf edge up to ~ 50km offshore. The area comprises important habitat for breeding aggregations of humpback whales (*Megaptera novaeangliae*), with mother–calf pairs showing a strong preference for shallow waters within Antongil Bay. The IMMA also hosts migrating humpback whales, with demonstrated connectivity to several other breeding grounds and migration routes, including those in the Mascarenes, northwest Madagascar, southern Madagascar, the Comoros Archipelago, and eastern Africa. Diversity and densities of other species within the area are observed to be relatively moderate for the region; however, Vulnerable sperm whales (*Physeter macrocephalus*) and beaked whale species have been observed in association with the 2000 m isobath.

Rosenbaum, 2003; Cerchio et al., 2009; Best and Brandão, 2009; Cerchio et al., 2016; Trudelle et al. 2016; Dulau et al., 2017; Trudelle et al., 2018; Fig. 1). In addition to humpback whales, at least eight other species were sighted during the REMMOA aerial survey in northeast Madagascar (January 2010). Of note, a number of deep-diving species were observed to associate with the 2000 m depth contour, including vulnerable sperm whales, near-threatened false killer whales, Cuvier's beaked whales, Longman's beaked whales, and other *Mesoplodon* spp. (Van Canneyt et al., 2010). Sperm whale relative density (not corrected for availability bias) was calculated as 0.20×10^{-2} individuals per km^2 , and uncorrected beaked whale relative density was calculated as 0.25×10^{-2} individuals per km^2 (Laran et al., 2012, 2017).

Criterion B: Distribution and Abundance

Sub-criterion B2: Aggregations

The area represents an important aggregation area for breeding humpback whales. Kernel density models of satellite telemetry data from whales initially tagged off Reunion Island indicate breeding hotspots off Île St. Marie, including the Saint Marie Channel and the waters of the continental shelf south of Île Sainte Marie to the Mongoro River mouth (Dulau et al., 2017).




Figure 1: Distribution of the humpback whale sightings made in the Sainte Marie Channel during the breeding season (June-September) across the five-year study period (2009-2013). Each dot represents a sighting. Adapted from Trudelle et al., 2018, Fig. 1.

Similarly, Cerchio et al. (2016) identified the same general area as an area of "localized" movement behaviour for whales tagged off Île Sainte-Marie in 2012. These 'hotspots' coincide with a relatively shallow area of the continental shelf, which may provide a higher suitability of breeding habitat for this species (Trudelle et al., 2016). While the sample size of satellite tagged animals is relatively small compared to the size of the population (although comparative with other satellite telemetry studies for this species), research suggests that humpback whales have a strong tendency to aggregate when conspecifics are detected (e.g., by song), especially now populations have been diminished by whaling (Clapham and Zerbini, 2015). As such, a breeding "hotspot" indicated by only a few animals through satellite telemetry data may represent a more significant aggregation area as other animals are attracted to the site.

Criterion C: Key Life Cycle Activities

Sub-Criterion C1: Reproductive Areas

The southwest Indian Ocean represents breeding and migratory habitat for humpback whale "Breeding Stock C," as defined by the International Whaling Commission. Breeding Stock C is further divided into four "breeding substocks" (i.e., "BSC1-C4"). Breeding Substocks C3 and C4 are known to aggregate in the coastal waters of central eastern Madagascar, including Antongil Bay, Île Sainte-Marie, and the waters south to the Mongoro River mouth during the winter (Rosenbaum et al., 1997; Best et al., 1998; Ersts and Rosenbaum, 2003; Cerchio et al. 2009; Best and Brandão, 2009; Cerchio et al., 2016; Trudelle et al., 2016; Dulau et al., 2017; Trudelle et al., 2018). Following preliminary surveys in 1996, Rosenbaum et al. (1997) described Antongil Bay as a breeding area for humpback whales in the southwestern Indian Ocean. Behaviours widely accepted to indicate breeding activity were regularly sighted, including males singing and exhibiting competitive behaviour, and females with young calves (Pomilla and Rosenbaum, 2006). Humpback whales have generally been observed in Antongil Bay from June to October, with the highest concentrations occurring in July through early September (Rosenbaum et al.,




Figure 2: Tracks of all tagged humpback whales (A) from Île Sainte-Marie in the northeast during 2012, with (B) detail of movements off the central east coast, and (C) from Anakao in the southwest during 2013, with (D) detail of movements off the southwest to southeast coasts. Each track is represented by dots for speed-filtered locations and a line connecting temporally consecutive locations. Females are represented in yellow, males in red, and a single whale of unknown sex in green. The satellite tracks clearly demonstrate the importance of this region, including the IMMA, as migratory habitat for humpback whales (Criterion C3). Adapted from Cerchio et al., 2016, Figure 2.

1997; Cerchio et al., 2009). Photographic mark-recapture data indicated that humpback whales utilizing Antongil Bay between 2000 and 2006 had low recapture rates with short residency times; therefore, whales appeared to regularly move through the bay with transient residency as part of a more extensive breeding range for a large population, estimated to be in excess of 6000 animals in 2006 (Cerchio et al., 2009). More recently, satellite telemetry data of humpback whales tagged off Île Sainte-Marie (N=12) in 2012, Anakao in the southwest (N=11), and La Reunion (N=12) in 2013 have highlighted potential breeding 'hotspots' off Île Sainte-Marie and the waters south to the Mongoro River mouth, but no entry or usage of Antongil Bay despite substantial use of the Sainte Marie Channel and areas just east of the Bay mouth (Cerchio et al., 2016; Trudelle et al., 2016; Dulau et al., 2017; Fig. 2). These data and other anecdotal observations suggest a reduced usage of Antongil Bay in recent years, and potential change of condition given extensive documentation of high densities in the bay during peak season in previous years (1996 to 2006) (Rosenbaum et al., 1997, Ersts and Rosenbaum, 2003, Cerchio et al., 2009).

Criterion C: Key Life Cycle Activities Sub-Criterion C3: Migration Routes

There are multiple sources of evidence of extensive humpback whale movements between the area defined by the Madagascar Central East Coast IMMA and other areas surrounding Madagascar, as well as areas off east Africa and the Mascarenes (e.g., Rosenbaum et al., 2009; Ersts et al., 2011; Fossette et al., 2014; Cerchio et al., 2008, 2016; Trudelle et al., 2016; Dulau et al., 2017; Kershaw et al., 2017) (Fig. 2). More broadly, the area falls within the migratory stream of Breeding Stock C, as animals transit annually between productive feeding areas in the Southern Ocean and warmer wintering grounds off East Africa, the Comoros Archipelago, Madagascar, and the Mascarenes.

Supporting Information

Best, P.B., Findlay, K.P., Sekiguchi, K., Peddemors, V.M., Rakotonirina, B., Rossouw, A., Gove, D. 1998. 'Winter distribution and possible migration routes of humpback whales *Megaptera novaeangliae* in the southwest Indian Ocean'. Marine Ecology Progress Series 162: 287-299.

- Best PB, Brandão A. 2009. Humpback whaling at Madagascar, 1910–1950. International Whaling Commission document SCF09SH2, 22–25 June 2009, Madeira, Portugal.
- Cerchio S, Findlay K, Ersts P, Minton G, Bennet D, Meyer M et al. 2008. Initial assessment of exchange between breeding stocks C1 and C3 of humpback whales in the western Indian Ocean using photographic mark–recapture data, 2000–2006. International Whaling Commission document SC/60/SH33, 23–27 June 2008, Santiago, Chile.
- Cerchio S, Ersts P, Pomilla C, Loo J, Razafindrakoto Y, Leslie M et al. 2009. Updated estimates of abundance for humpback whale breeding stock C3 off Madagascar, 2000–2006. International Whaling Commission document SC/61/SH7, 22–25 June 2009, Madeira, Portugal.
- Cerchio S, Trudelle L, Zerbini AN, Charrassin JB, Geyer Y, Mayer FX et al. 2016. Satellite telemetry of humpback whales off Madagascar reveals insights on breeding behavior and long-range movements within the southwest Indian Ocean. *Marine Ecology Progress Series* 562: 193–209.
- Clapham, P.J. and Zerbini, A.N., 2015. Are social aggregation and temporary immigration driving high rates of increase in some Southern Hemisphere humpback whale populations? *Marine biology*, 162(3), pp.625–634.
- Dulau V, Pinet P, Geyer Y, Fayon J, Mongin P, Cottarel G et al. 2017. Continuous movement behavior of humpback whales during the breeding season in the southwest Indian Ocean: On the road again! *Movement Ecology* 5: 11.
- Ersts PJ, Rosenbaum HC. 2003. Habitat preference reflects social organization of humpback whales (*Megaptera novaeangliae*) on a wintering ground. *Journal of Zoology* 260: 337–345.
- Ersts PJ, Pomilla C, Kiszka J, Cerchio S, Rosenbaum HC, Vély M, et al. 2011. Observations of individual humpback whales utilizing multiple migratory destinations in the south-western Indian Ocean. *African Journal of Marine Science* 33: 333–338.
- Faria MA, DeWeerd, J, Pace F, Mayer FX. 2013. Observation of a humpback whale (*Megaptera novaeangliae*) birth in the coastal waters of Sainte Marie Island, Madagascar. *Aquatic Mammals* 39: 296–305.
- Fossette S, Heide-Jørgensen MP, Jensen MV, Kiszka J, Bérubé M, Bertrand N, Vély M. 2014. Humpback whale (*Megaptera novaeangliae*) post-breeding dispersal and southward migration in the western Indian Ocean. *Journal of Experimental Marine Biology and Ecology* 450: 6–14
- Kershaw F, Carvalho I, Loo J, Pomilla C, Best PB, Findlay KP, et al. 2017. Multiple processes drive genetic structure of humpback whale (*Megaptera novaeangliae*) populations across spatial scales. *Molecular ecology*. 26: 977–94.
- Laran, S., Authier, M., Van Canneyt, O., Doremus, G., Watremez, P., Ridoux, V. (2017). A Comprehensive Survey of Pelagic Megafauna: Their Distribution, Densities, and Taxonomic Richness in the Tropical Southwest Indian Ocean. *Front. Mar. Sci.* 4:139. doi: 10.3389/fmars.2017.00139
- Laran, S., Van Canneyt, O., Dorémus, G., Mannocci, L., Ridoux, V. & Watremez, P. 2012. Distribution et abondance de la mégafaune marine dans le sud-ouest de l'océan Indien tropical. REMMOA-Océan Indien. Rapport final pour l'Agence des Aires Marines Protégées. 168p.
- Pomilla, C., and Rosenbaum, H.C. 2006. Estimates of relatedness in groups of humpback whales (*Megaptera novaeangliae*) on two wintering grounds of the Southern Hemisphere. *Molecular Ecology* 15, 2541–2555.

Rosenbaum HC, Walsh P, Razafindrakoto Y, Vely M, Desalle R. 1997. First description of a humpback whale wintering ground in Baie d'Antongil, Madagascar. *Conservation Biology* 11: 308–314.

Rosenbaum, H.C., Pomilla, C., Mendez, M., Leslie, M.S., Best, P.B., Findlay, K.P., Minton, G., Ersts, P.J., Collins, T., Engel, M.H. and Bonatto, S.L., 2009. Population structure of humpback whales from their breeding grounds in the South Atlantic and Indian Oceans. *PLoS One*, 4(10), p.e7318.

Trudelle, L., Cerchio, S., Zerbini, A.N., Geyer, Y., Mayer, F.X., Jung, J.L., Hervé, M.R., Pous, S., Sallée, J.B., Rosenbaum, H.C. and Adam, O., 2016. Influence of environmental parameters on movements and habitat utilization of humpback whales (*Megaptera novaeangliae*) in the Madagascar breeding ground. *Royal Society open science*, 3(12), p.160616.

Trudelle L, Charrassin J-B, Saloma A, Pous S, Kretzschmar A, and Adam O. 2018. First insights on spatial and temporal distribution patterns of humpback whales in the breeding ground at Sainte Marie Channel, Madagascar. *African Journal of Marine Science* 40: 75-86.

Van Canneyt, O., Dorémus, G., Laran, S., Ridoux, V., Watremez, P., 2010. REMMOA Sud Ouest Océan Indien: Rapport intermédiaire pour l'Agence des Aires Marines Protégées, 70 pp.

Acknowledgements

We would like to thank the participants of the 2019 IMMA Regional Expert Workshop held in Salalah, Oman for the identification of IMMAs in the Western Indian Ocean and Arabian Seas. Funding for the identification of this IMMA was provided to the Global Ocean Biodiversity Initiative by the International Climate Initiative (IKI). The German Federal Ministry for the Environment, Nature Conservation and Nuclear Safety (BMU) supports this initiative on the basis of a decision adopted by the German Bundestag. Support was also provided by Whale and Dolphin Conservation and the Tethys Research Institute.



**MARINE MAMMAL
PROTECTED AREAS
TASK FORCE**



IMMA

Supported by:



Federal Ministry for the
Environment, Nature Conservation,
Building and Nuclear Safety



based on a decision of the German Bundestag

Suggested Citation: IUCN-Marine Mammal Protected Areas Task Force, 2021. Madagascar Central East Coast IMMA Factsheet. <https://www.marinemammalhabitat.org/wp-content/uploads/imma-factsheets/WesternIndianOcean/madagascar-central-east-coast-WesternIndianOcean.pdf>. Downloaded on (day month year).

PDF made available for download at
<https://www.marinemammalhabitat.org/wp-content/uploads/imma-factsheets/WesternIndianOcean/madagascar-central-east-coast-WesternIndianOcean.pdf>