

## Area Size

120,614 km<sup>2</sup>

## Qualifying Species and Criteria

Omura's whale – *Balaenoptera omurai*  
Criteria B1

Antarctic blue whale – *Balaenoptera musculus intermedia*; Criteria A

Pygmy blue whale – *Balaenoptera musculus brevicauda*; Criterion A, C3

Fin whale – *Balaenoptera physalus*  
Criterion A

Humpback whale – *Megaptera novaeangliae*  
Criteria B2, C1, C3

Indian Ocean humpback dolphin – *Sousa plumbea*  
Criteria A, B1

Melon-headed whale – *Peponocephala electra*  
Criterion B2

Dugong – *Dugong dugon*  
Criterion A

Sperm whale – *Physeter macrocephalus*  
Criterion A

# Northwest Madagascar and Northeast Mozambique Channel IMMA

## Marine Mammal Diversity (D2)

*Balaenoptera acutorostrata*, *Balaenoptera bonaerensis*, *Balaenoptera musculus brevicauda*, *Balaenoptera musculus intermedia*, *Balaenoptera omurai*, *Balaenoptera physalus*, *Dugong dugon*, *Globicephala macrorhynchus*, *Grampus griseus*, *Indopacetus pacificus*, *Kogia sima*, *Lagenodelphis hosei*, *Megaptera novaeangliae*, *Peponocephala electra*, *Physeter macrocephalus*, *Pseudorca crassidens*, *Sousa plumbea*, *Stenella attenuata*, *Stenella longirostris*, *Tursiops aduncus*, *Tursiops truncatus*, *Ziphius cavirostris*

## Summary

The waters of the Mozambique channel, off the northwest coast of Madagascar, comprise a highly diverse and important region for marine mammals. A minimum of 23 marine mammal species have been documented here, including 6 baleen whale species, 16 toothed whales and dolphins, and the dugong. The area encompasses important habitats for breeding, feeding and migration, as well as small and resident populations for several species. Species listed in the IUCN Red List of Threatened Species include: Critically Endangered Antarctic blue whales (*Balaenoptera musculus intermedia*); Endangered pygmy blue whales (*B.m. brevicauda*); Endangered Indian Ocean humpback dolphins (*Sousa plumbea*); Vulnerable fin whales (*Balaenoptera physalus*), Vulnerable sperm whales (*Physeter macrocephalus*), and Vulnerable dugongs. Notably, a population of Omura's whales is resident in the area year-round and is likely regionally isolated. During a large-scale aerial survey, encounter rates for several species, including common dolphins (*Delphinus delphis*) and Indo Pacific bottlenose dolphins (*Tursiops aduncus*), melon-headed whales (*Peponocephala electra*), and spinner or spotted dolphins (*Stenella sp.*) were particularly high on the shelf and slope habitats of the IMMA.

## Description

This IMMA encompasses a stretch of the west coast of Madagascar from ca. 12°S to 16°S offshore to approximately 100 km. It encompasses a large variety of habitats, from coastal mangroves, coral reefs and sandy and rocky shores, to neritic continental shelf and pelagic deep-water habitats. The area includes the Nosy Be region from approximately 12.7°S to 13.7°S, including the islands of Nosy Be, Nosy Komba, Nosy Iranja and the Nosy Mitsio archipelago, where relatively extensive research on marine mammals has been conducted.

At minimum, 23 marine mammal taxa (22 species and 1 subspecies) have been documented in this area (Van Canneyt et al., 2010; Cerchio et al., in press). The IMMA encompasses important habitat in three specific zones, each with a distinct species, or species group, that satisfies a range of IMMA criteria. The habitat zones and associated species are as follows. The near-coastal zone: Indian Ocean humpback dolphin, Indo-Pacific bottlenose dolphin and dugong. The continental shelf zone: Omura's whale, humpback whale, dwarf minke whale, spinner dolphin, pantropical spotted dolphin, and Indo-Pacific bottlenose dolphin. And the offshore continental slope and abyssal plain deep-water zone: Antarctic blue whale, pygmy blue whale, fin whale, Antarctic minke whale, spinner dolphin, pantropical spotted dolphin, Fraser's dolphin, common bottlenose dolphin, Risso's dolphin, short-finned pilot whale, melon-headed whale, killer whale, false killer whale, sperm whale, Cuvier's beaked whale, Longman's beaked whale, dwarf and pygmy sperm whale.

Much of the coastline is remote and undeveloped, yet there are several highly-developed and heavily-populated centres, including the port of Mahajanga and the tourism hubs of the Nosy Be region and Antsiranana. Bycatch of small cetaceans has been documented in coastal artisanal fisheries and although hunting appears to be much less prevalent than it is in the southwest region, it is not entirely absent (Cerchio et al. 2014; 2015b). Currently there is extensive seismic exploration planned throughout this entire region which presents a distinct threat to sensitive populations of cetaceans (Cerchio et al., 2019). There is also coastal industrial development

in the form of mining for Rare Earth Elements that has the potential to pollute coastal waters in the Nosy Be region (Cerchio et al., 2019).

## Criterion A: Species or Population Vulnerability

Indian Ocean humpback dolphin, *Sousa plumbea* (IUCN Red List EN, Braulik et al., 2015; 2017). The conservation status of the Indian Ocean humpback dolphin is reported to be heterogeneous along the west coast of Madagascar; population(s) in the southwest are highly impacted (where active hunting occurs in the waters of the Vezo people), whereas population(s) in the northwest appear to be healthier and more abundant (Cerchio et al., 2009; 2014; 2015b; Razafindrakoto, 2004). The regions encompassed by the IMMA, well documented around the Nosy Be, Nosy Iranja and Nosy Mistio area, appear to be a haven for humpback dolphins relative to much of the other parts of their range in Madagascar.

Antarctic blue whale, *Balaenoptera musculus intermedia* (IUCN Red List CR, Cooke 2018a) and SWIO pygmy blue whale, *Balaenoptera musculus brevicauda* (IUCN Red List EN, Cooke 2018b). Passive acoustic monitoring from recording sites on the shelf break near Nosy Be indicated the presence of both species off northwest Madagascar (Cerchio et al.,) 2018b. Some acoustic detections were high signal-to-noise ratio indicating presence within 10km of the shelf break, whereas it was suspected that most detections were within 100km of the coast. The offshore extent of the IMMA boundary was chosen to roughly encompass the likely range at which blue whales were detected acoustically during the austral winter and fall migrations, but it is likely that whales range further offshore into the Mozambique Channel (Cerchio et al. 2018).

Sperm whales, *Physeter macrocephalus*, listed as *Vulnerable* (Taylor et al. 2008) have not been a focus of study in the waters of northwest Madagascar, however they use the deep offshore waters of the IMMA regularly and were encountered over the slope habitat during REMMOA aerial survey. Visual surveys of deep waters in Nosy Be resulted in encounters near the 2000m depth contour, and acoustic monitoring produced numerous detections.

Dugong, *Dugong dugon*, IUCN Red List Vulnerable (Marsh and Sobotzick, 2015) is also encountered in the area (Cerchio et al. 2012; Van Canneyt et al., 2010). The species is thought to have been historically widespread throughout Madagascar, but believed to be heavily exploited to the point that sightings of live animals are very rare. Population status is generally uncertain and it is possible/likely to have been extirpated throughout much of its former range (Cooke et al. 2003; Cerchio et al., 2012; Davis et al., in press). Interview surveys with fishers along the west coast of Madagascar indicated decline in numbers of sightings and individuals captured in hunting and by-catch during the decade 2000-2010, and shifts in the relative distribution of reports over time, suggesting that populations in some locations may be more impacted or closer to extirpation (Cerchio et al. 2012).

## **Criterion B: Distribution and Abundance**

### **Sub-criterion B1: Small and Resident Populations**

Omura's whale, *Balaenoptera omurai* – Existing data, much of it recent (including year-long acoustic monitoring, visual surveys, photographic identification, and satellite telemetry), indicate that this is a resident, non-migratory population whose distribution is likely determined by local shallow water ecological processes and patchy and ephemeral prey resources. Furthermore, this population of Omura's whale may be isolated within a fragmented oceanic/global range for the species. The boundaries of the area encompass the range documented by boat surveys and satellite telemetry data including the offshore extent of documented movements, and likely most of the habitat for this population as indicated by habitat suitability modelling (Cerchio et al., 2015a; 2018a). Indian Ocean humpback dolphin, *Sousa plumbea*. The latitudinal range of the IMMA encompasses most, if not all, of the coastal habitat for this species within that area where hunting pressure is believed to be least in Madagascar (Cerchio et al., 2009, 2014, 2015b; Razafindrakoto 2004). Dugong, *Dugong dugon*- There are interview accounts of relatively recent sightings, bycatch and hunting in the northwest Nosy Be region (Cerchio et al. 2012) and in the Diana region in the extreme northwest (C3-MIOIP, 2010b; Davis et al., in press), and aerial survey documentation of live

sightings between Mahajanga and Bay of Sahamalaza in 2009 (Van Conneyt et al., 2010). These data suggest that the northwest of Madagascar may constitute a range where a remnant viable population remains, but is still under hunting and bycatch pressure and likely in decline. During the REMMOA aerial survey (between mid December 2009 and early January 2010) relative density of dugong was estimated at  $0.10 \times 10^{-2}$  individuals.km<sup>2</sup> (CV: 74%), corresponding to probably fewer than 200 dugongs in this area (Laran et al., 2012). The coastal boundaries of the area encompass all of these sites and the majority of the most recent sightings and documentation where a remnant viable population may remain in Madagascar (Cooke et al., 2003; C3-MIOIP, 2010; Van Canneyt et al., 2010; Cerchio et al., 2012; Davis et al., in press).

## **Criterion C: Key Life Cycle Activities**

### **Sub-criterion C1: Reproductive Areas**

Humpback whale, *Megaptera novaeangliae* – Madagascar is a well-documented breeding area for humpback whales, and small boat surveys indicate that they are common in coastal waters of the Nosy Be region during the mid to late breeding season; year-long acoustic monitoring in the deep offshore waters of the Nosy Be region indicates near continuous presence of humpback whale song from early July to mid-November (Cerchio et al. 2018b; in press).

Antarctic blue whale, *Balaenoptera musculus intermedia* – Year-long acoustic monitoring in the deep offshore waters off the Nosy Be region indicated the presence of Antarctic blue whale song off northwest Madagascar throughout the Austral winter from June to September, suggesting previously unrecognized breeding season habitat (Cerchio et al., 2018b). Fin whale, *Balaenoptera physalus* – Year-long acoustic monitoring in the deep offshore waters off the Nosy Be region indicates the presence of fin whales during the late Austral winter, from early August to mid-September (Cerchio et al. 2018b). The timing of fin whale song suggests a later arrival than Antarctic blue whales and a lower rate of occurrence and occupancy, suggesting the northern extent of breeding habitat.




Figure 1: Sightings of cetaceans collected on effort during REMMOA aerial survey in the North of Mozambique Channel (between mid December 2009 and early January 2010) detailed by species or taxa and encounter rate of individual (for cell with up to 50km of effort - Unpublish map Pelagis-AFB; Van Canneyt et al. 2010)

## Criterion C: Key Life Cycle Activities Sub-criterion C3: Migration Routes

Pygmy blue whale, *Balaenoptera musculus breviceuda* – Year-long acoustic monitoring in the deep offshore waters of the Nosy Be region indicates presence of the Madagascar-type pygmy blue whale song with bi-modal peaks of singing activity during May-July and October-January (Cerchio et al 2018b). This pattern suggests a previously unrecognized migratory corridor between summer feeding and winter breeding grounds south and north of Madagascar, respectively. This migratory corridor likely connects feeding habitat on the Madagascar Plateau and Ridge (Best et al. 2003) and likely breeding habitat off Kenya and the Seychelles area (Branch et al. 2007, Barber et al. 2016). Humpback whale, *Megaptera novaeangliae* – Satellite telemetry data indicates that this area of coast is a late season migratory corridor for humpback whales (Fossette et al 2014, Dulau et al 2017).

## Criterion D: Special Attributes Sub-criterion D2: Diversity

A minimum of 22 species of cetaceans and the dugong have been documented in the Nosy Be region (Van Canneyt et al. 2010; Cerchio et al., in press). In addition to the species described above for other criteria, the following species are included for this criterion. Dwarf minke whale song vocalizations were detected on several days during non-systematic review of acoustic data from continental shelf waters around Nosy Be; it is possible that future analyses will indicate regular presence during the Austral winter, and thus in breeding habitat (Cerchio et al., in press). Spinner dolphins are the most often sighted species by number of individuals in the continental slope and offshore areas (Cerchio et al., 2014). Pantropical spotted dolphins are frequently sighted in offshore areas during surveys around Nosy Be, at times in groups exceeding 500 individuals, and in combination with spinner dolphins, as well as on

the continental shelf in smaller-sized groups (Cerchio et al. 2014; in press). Fraser's dolphin were occasionally encountered in deep waters around Nosy Be, as were short-finned pilot whales, with groups of up to more than 40 individuals (Cerchio et al. 2014; in press). Melon-headed whale and false killer whale were also occasionally encountered in shelf waters by tourists (Cerchio, unpublished data). During the REMMOA aerial survey, bottlenose dolphins (unclear if common or Indo-Pacific), have a relative density of  $7 \times 10^{-2}$  individuals.km<sup>2</sup> (CV: 30%) over the shelf area of Comoros and the Malagasy coast (Laran et al., 2012) while *Stenella spp.* dominate the community with a density peaking to more than  $30 \times 10^{-2}$  individuals.km<sup>2</sup> (CV: 59%) over the slope strata (Laran et al, 2012). Nevertheless, most of the large Delphinidae (*Tursiops spp.*) were encountered along the Malagasy coast, while *Stenella spp.* concentrated around the Comoros archipelago. Melon-headed whales, encountered in large groups resulting in a density of  $50 \times 10^{-2}$  individuals.km<sup>2</sup> (CV: 42%) for small blackfish (Globicephalinae) over the slope of this area and around the Comoros islands. The slope area has also sperm whale, beaked whales, Cuvier's and Longman's beaked whale, false killer whale, short-finned pilot whale and *Kogia spp.* (Van Canneyt et al., 2010; Cerchio et al., in press).

## Supporting Information

Barber, R., Sikora, I. and Nimak-Wood, M., 2016. Blue whales *Balaenoptera musculus* in offshore waters of Kenya. *African Journal of Marine Science*, 38(2):279-284.

Best, P.B., Rademeyer, R.A., Burton, C., et al. 2003. The abundance of blue whales on the Madagascar Plateau, December 1996. *Journal of Cetacean Research and Management*, 5: 253–260.

Branch, T.A., Stafford, K.M., Palacios, D.M., et al. 2007. Past and present distribution, densities and movements of blue whales *Balaenoptera musculus* in the Southern Hemisphere and northern Indian Ocean. *Mammal Review*, 37:116–175.

Braulik, G.T., Findlay, K., Cerchio, S., Baldwin, R., Perrin, W. 2017. *Sousa plumbea*. The IUCN Red List of Threatened Species 2017: e.T82031633A82031644. <http://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T82031633A82031644.en>.

Braulik, G. T., Findlay, K., Cerchio, S. and Baldwin, R. 2015. Assessment of the Conservation Status of the Indian Ocean Humpback Dolphin (*Sousa plumbea*) Using the IUCN Red List Criteria. In: Jefferson T., and Curry B. (Eds). *Humpback Dolphins (Sousa spp.): Their Current Status and Conservation*, pPp. 119-141. *Advances in Marine Biology*, 72. Oxford: Academic Press.

C3 Madagascar and Indian Ocean Islands Programme, 2010b. Rapid Assessment of Dugongs and their Seagrass Habitat in Northern Madagascar. A Report Submitted to the Convention On Migratory Species Of Wild Animals (CMS) Secretariat, Abu Dhabi. C3 Technical Report Series No. 6. ISSN 1754-5188. Community Centered Conservation (C3), London, UK. 19pp.

Cerchio, S., Laran, S., Andrianarivelo, N., Saloma, A., Andriananteniana, B., Van Canneyt, O., and Rasoloarijao, T. In press. Cetacean Species Diversity in Malagasy Waters. In: Goodman, S. M. (ed). *The New Natural History of Madagascar*. Princeton: Princeton University Press.

Cerchio, S., Yamada, T.K., Brownell, Jr. RL. 2019. Global distribution of Omura's whales (*Balaenoptera omurai*) and assessment of range-wide threats. *Frontiers in Marine Ecology* 6:67. doi: 10.3389/fmars.2019.00067.

Cerchio, S., and Yamada, T. 2018. 'Omura's whale: *Balaenoptera omurai*'. In: Würsig B, Thewissen JGM, and Kovacs K (eds) "Encyclopedia of Marine Mammals", pp. 656-659, 3rd ed., Academic Press, San Diego, CA, USA.

Cerchio, S, Andrianantenaina, B., Zerbini, A., Pendleton, D., Rasoloarijao, T., Cholewiak, D. 2018a. Residency, feeding ecology, local movements and potential isolation of the Madagascar Omura's whale (*Balaenoptera omurai*) population. Paper SC/67B/NH09 presented to the IWC Scientific Committee.

- Cerchio, S, Rasoloarijao, T, Cholewiak, D. 2018b. Acoustic monitoring of Blue Whales (*Balaenoptera musculus*) and other baleen whales in the Mozambique Channel off the Northwest Coast of Madagascar. Paper SC/67B/SH/14 presented to IWC Scientific Committee.
- Cerchio, S, Andrianantenaina, B, Lindsay, A, Rekdahl, M, Andrianarivelo, N, and Raoloarijao, T. 2015a. Omura's whales (*Balaenoptera omurai*) in Northwest Madagascar: ecology, behavior and conservation needs. Royal Society Open Science 2: 150301. DOI:10.1098/rsos.150301.
- Cerchio, S, Andrianarivelo, N, and Andrianantenaina, B. 2015b. 'Ecology and conservation status of Indian Ocean humpback dolphins (*Sousa plumbea*) in Madagascar'. In: Jefferson T, and Curry B (Eds). Humpback Dolphins (*Sousa* spp.): Their Current Status and Conservation, pp. 163-199. Advances in Marine Biology Series, 72. Oxford: Academic Press.
- Cerchio, S, Andrianarivelo, N, Andrianantenaina, B, Cordi, V. 2014. Ecology, status, fisheries interactions and conservation of coastal Indian Ocean humpback dolphins and Indo-Pacific bottlenose dolphins on the west coast of Madagascar. Paper SC/65B/SM21 presented to the IWC Scientific Committee.
- Cerchio, S, Andrianarivelo, N, Andrianantenaina, B. 2012a. Sightings, incidental by-catch, and hunting of dugongs (*Dugong dugon*) on the west coast of Madagascar, as Inferred from Interviews with Local Fishers. Report to the Convention on Migratory Species. 17pp.
- Cerchio, S, Andrianarivelo, N, Razafindrakoto, Y, Mendez, M, and Rosenbaum, H. 2009. Coastal dolphins hunting in the southwest of Madagascar: status of populations, human impacts and conservation actions. Paper SC/61/SM15 presented to the IWC Scientific Committee.
- Cooke, A., Lutjeharms, J.R.E., and Vasseur, P. 2003. 'Marine and Coastal Ecosystems'. In: Goodman, S. and Benstead, J. (eds) Natural History of Madagascar, p 179-209. Chicago University Press, Chicago, IL.
- Cooke, J.G. 2018a. *Balaenoptera musculus* ssp. *intermedia*. The IUCN Red List of Threatened Species 2018: e.T41713A50226962. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T41713A50226962.en>. Downloaded on 25 February 2019.
- Cooke, J.G. 2018b. *Balaenoptera musculus*. The IUCN Red List of Threatened Species 2018: e.T2477A50226195. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T2477A50226195.en>. Downloaded on 24 February 2019.
- Davis, P. Z. R., Cervchio, S., Andrianarivelo, N. In press. *Dugong dugon*, *Dugong lambohara*. In: Goodman, S. M. (ed). The New Natural History of Madagascar. Princeton: Princetown University Press.
- Dulau, V, Pinet, P, Geyer, Y, Fayan, J, Mongin, P, Cottarel, G, Zerbini, A, Cerchio, S. 2017. Movement behavior of humpback whales during the breeding season: on the road again! Movement Ecology 5:11. DOI 10.1186/s40462-017-0101-5.
- Fossette, S, Heide-Jørgensen, MP, Jensen, MV, Kiszka, J, Bérubé, M, Bertrand, N, Vély, M 2014. Humpback whale (*Megaptera novaeangliae*) post breeding dispersal and southward migration in the western Indian Ocean. Journal of Experimental Marine Biology and Ecology, 450: 6-14
- Laran, S., Van Canneyt, O., Dorémus, G., Mannocci, L., Ridoux, V. & Watremez, P. 2012. Distribution et abondance de la mégafaune marine dans le sud-ouest de l'océan Indien tropical. REMMOA-Océan Indien. Rapport final pour l'Agence des Aires Marines Protégées. (Distribution and abundance of the marine megafauna of the SouthWest Indian Ocean. Report for the Marine Protected Area Agency), 168p
- Laran, S., Authier, M., Van Canneyt, O., Doremus, G., Watremez, P., Ridoux, V. 2017. A Comprehensive Survey of Pelagic Megafauna: Their Distribution, Densities, and Taxonomic Richness in the Tropical Southwest Indian Ocean. Frontiers in Marine Science. 4:139. doi: 10.3389/fmars.2017.00139

Marsh, H. and Sobtzick, S. 2015. Dugong dugon. The IUCN Red List of Threatened Species 2015: e.T6909A43792211. <http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T6909A43792211.en>. Downloaded on 24 February 2019.

McDonald, MA, Hildebrand, JA, Mesnick, SL. 2006. Biogeographic characterization of blue whale song worldwide: using song to identify populations. *Journal of Cetacean Research and Management*, 8, 55–65.

Razafindrakoto, Y., Andrianarivelo, N., Rosenbaum, H.C., 2004. Sightings, catches, and other records of Indo-Pacific humpback dolphins in the coastal waters of Madagascar. *Aquatic Mammals*, 30: 103–110.

Taylor, B.L., Baird, R., Barlow, J., Dawson, S.M., Ford, J., Mead, J.G., Notarbartolo di Sciarra, G., Wade, P. and Pitman, R.L. 2008. *Physeter macrocephalus*. The IUCN Red List of Threatened Species 2008: e.T41755A10554884. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T41755A10554884.en>.

Van Canneyt, O., Dorémus, G., Laran, S., Ridoux, V. and Watremez, P. 2010. REMMOA Sud Ouest Océan Indien: Rapport intermédiaire pour l'Agence des Aires Marines Protégées (REMMOA SouthWest Indian Ocean, survey report for the Marine Protected Area Agency), 70 pp.

## Acknowledgements

We would like to thank the participants of the 2019 IMMA Regional Expert Workshop held in Salalah, Oman for the identification of IMMAs in the Western Indian Ocean and Arabian Seas. Funding for the identification of this IMMA was provided to the Global Ocean Biodiversity Initiative by the International Climate Initiative (IKI). The German Federal Ministry for the Environment, Nature Conservation and Nuclear Safety (BMU) supports this initiative on the basis of a decision adopted by the German Bundestag. Support was also provided by Whale and Dolphin Conservation and the Tethys Research Institute.



**MARINE MAMMAL  
PROTECTED AREAS  
TASK FORCE**



Supported by:



Federal Ministry for the  
Environment, Nature Conservation,  
Building and Nuclear Safety



based on a decision of the German Bundestag



TETHYS  
since 1986



WHALE AND  
DOLPHIN  
CONSERVATION

Suggested Citation: IUCN-Marine Mammal Protected Areas Task Force, 2021. Northwest Madagascar and Northeast Mozambique Channel IMMA Factsheet. <https://www.marinemammalhabitat.org/wp-content/uploads/imma-factsheets/WesternIndianOcean/northwest-madagascar-northeast-mozambique-channel-WesternIndianOcean.pdf>. Downloaded on (day month year).

PDF made available for download at <https://www.marinemammalhabitat.org/wp-content/uploads/imma-factsheets/WesternIndianOcean/northwest-madagascar-northeast-mozambique-channel-WesternIndianOcean.pdf>.